



4D-2: Problems with Logarithmic Functions

In this lesson we will use the properties of logarithms to solve equations that involve logarithms. We first need to recall these properties from earlier:

Properties of Logarithms

Let b , R , and S be positive real numbers with $b \neq 1$, and c a

Product Rule: $\log_b(PQ) = \log_b P + \log_b Q$

Quotient Rule: $\log_b \frac{P}{Q} = \log_b P - \log_b Q$

Power Rule: $\log_b P^c = c \log_b P$

Where b , R , and S be positive real numbers with $b \neq 1$, and c is any real number.

Example Assume that x and y are positive below.

a) Write as a sum of logarithms with no exponents: $\log \frac{3x^2}{y}$

$$\log 3 + 2 \log x - \log y$$

b) Write as a single logarithm: $3 \ln 2 - 2 \ln 4 + \frac{1}{2} \ln 16$

$$\ln 2^3 - \ln 4^2 + \ln 16^{1/2} = \ln \left(\frac{8}{16} \cdot 4 \right) = \ln 2$$

To solve equations with logarithms, we can do one of the following:

1. Set the equation equal to zero, graph the corresponding function, and find the zeros.
2. Write both sides of the equation as one logarithm with the same base, convert to exponential form, and solve.

After solving, you must check your domain to be sure that x is in the domain of the original function.

Example.Solve.

a) $\log_3(x + 1) = 4$

$$x = 80$$

b) $2 \log x + 3 \log 2 = \log 16$

$$\log(x^2 \cdot 2^3) = \log 16$$

$$8x^2 = 16$$

$$x^2 = 2$$

$$x = \pm\sqrt{2}$$

x cannot be negative in original problem! So the only solution is $x = \sqrt{2}$

Exercises

Assuming x and y are positive, use the properties of logarithms to write the expression as a sum or difference of logarithms or multiples of logarithms.

$$1. \log_2 y^5 \quad \mathbf{5 \log_2 y}$$

$$2. \log_2 \frac{2x^3}{y^2} \\ \log_2 2x^3 - \log_2 y^2 \\ \log_2 2 + 3 \log_2 x - 2 \log_2 y \\ \mathbf{1 + 3 \log_2 x - 2 \log_2 y}$$

$$3. \log 1000x^4 \\ \log 1000 + \log x^4 \\ \mathbf{3 + 4 \log x}$$

Assuming x , y , and z are positive, use properties of logarithms to write the expression as a single logarithm.

$$4. \ln y - \ln 3 \quad \mathbf{\ln \frac{y}{3}}$$

$$5. 4 \log y - \log z \quad \mathbf{\log \frac{y^4}{z}}$$

$$6. 3 \ln 2 - 2 \ln 4 \\ \ln \frac{2^3}{4^2} = \mathbf{\ln \frac{1}{2}} \\ \text{or ...} \\ 3 \ln 2 - 2 \ln 2^2 \\ 3 \ln 2 - 4 \ln 2 \\ \mathbf{- \ln 2}$$

Find the exact solution algebraically, obtain a numerical approximation, and check it by substituting into the original equation.

$$7. \log_4(1-x) = 1 \quad \mathbf{x = -3}$$

$$8. 3 \ln(x-3) + 4 = 5 \quad \mathbf{x = \sqrt[3]{e} + 3}$$

$$9. 3 - \log(x+2) = 5 \quad \mathbf{x = -\frac{199}{100}}$$

$$10. \frac{1}{2} \ln(x+3) - \ln x = 0 \\ \ln(x+3)^{\frac{1}{2}} = \ln x \\ (x+3)^{\frac{1}{2}} = x \\ x+3 = x^2 \\ x = \frac{1 \pm \sqrt{13}}{2}, \text{ by domain restrictions } \mathbf{x = \frac{1 + \sqrt{13}}{2}}$$

$$11. \log x - \frac{1}{2} \log(x+4) = 1 \\ \log \frac{x}{\sqrt{x+4}} = 1 \\ \frac{x}{\sqrt{x+4}} = 10 \\ x = 10\sqrt{x+4} \\ x^2 = 100(x+4) \\ x^2 - 100x - 400 = 0 \\ x = 50 - 10\sqrt{29} \text{ or } x = 10\sqrt{29} + 50$$

$$\mathbf{\text{Solution: } x = 10\sqrt{29} + 50}$$

$$13. \log(x-2) + \log(x+5) = 2 \log 3 \\ \log(x-2)(x+5) = \log 3^2 \\ x^2 + 3x - 10 = 9 \\ x^2 + 3x - 19 = 0 \\ x = \frac{-\sqrt{85} - 3}{2} \text{ or } x = \frac{\sqrt{85} - 3}{2}$$

$$\text{Solution: } \mathbf{x = \frac{-3 + \sqrt{85}}{2}}$$

$$12. \ln(x-3) + \ln(x+4) = 3 \ln 2 \\ \ln(x-3)(x+4) = \ln 2^3 \\ x^2 + x - 12 = 8$$

$$x = -5 \text{ or } x = 4$$

$$\mathbf{\text{so ... } x = 4}$$

14. Determine whether a linear, logarithmic, exponential, power, or logistic regression equation is the best model for the data using your calculator and finding the R^2 value.

Year	1900	1910	1920	1930	1940	1950	1960	1970	1980	1990	2000
Alaska's Population	63.6	64.4	55.0	59.2	72.5	128.6	226.2	302.6	401.9	550.0	626.9

Best Curve = Logistic

15. In Chemistry, pH value is determined by the hydrogen ion concentration (H^+) relative to pure water. The function for finding pH is $p(H^+) = 7 + \log\left(\frac{1}{H^+}\right)$.
- State the domain of the function $p(H^+)$.
 - For lemon juice, we know $H^+ \approx 100,000$. Find the pH of lemon Juice.
 - If a solution of bleach has a pH value of $p(H^+) = 13.2$, find the value of H^+ .